

Ordinary Differential Equations of the Probability Functions of the Weibull Distribution and their Application in Ecology

H. 1. Okagbue¹, M. O. Adamu² and T. A. Anake³

^{1,2,3}Department of Mathematics, Covenant University, Ota, Ogun State, Nigeria

²Department of Mathematics, University of Lagos, Lagos State, Nigeria

Email : ¹hilary.okagbue@covenantuniversity.edu.ng

ABSTRACT

Weibull distribution has been applied to many areas in ecological studies and engineering. Application of the Weibull and other probability distributions in ecology are mainly in fitting ecological data which is very vital in revealing latent characteristics of the object of study. The use of the ordinary differential equations (ODE) in fitting has not been studied in ecological studies. Ordinary differential calculus was used to obtain the homogenous ODE of the probability density function (PDF), quantile function (QF), survival function (SF), inverse survival function (ISF), hazard function (HF) and reversed hazard function (RHF) whose solutions are their respective functions of the Weibull distribution. Different classes of ODEs were obtained. The novelty of this proposed method is applied to radiation data.

Keywords: Weibull distribution, Ordinary differential equations, probability function, data fit, parameters.

Mathematics Subject Classification: 46N30, 60E05, 62E10, 60E10, 60E99

Computing Classification System : G.3

Journal of Economic Literature (JEL) Classification: Q12, D24

1. INTRODUCTION

Calculus is a very key tool in the determination of mode of a given probability distribution and in estimation of parameters of probability distributions, amongst other uses. Probability density function (PDF) can be expressed as ODE whose solution is the PDF. Some of which are available. They include: Laplace distribution (Johnson et al., 1994), beta distribution (Elderton, 1906), raised cosine distribution (Rinne, 2010), Lomax distribution (Balakrishnan and Lai, 2009), eta prime distribution or Inverted beta distribution (Johnson et al., 1995).

1.1 Weibull Distribution in Ecological Studies

Weibull distribution has been applied to many areas in ecological studies. Application of the Weibull and other probability distributions in ecology are mainly in fitting ecological data which is very vital in revealing latent characteristics of the object of study.

Stauffer (1979) derived in details, the mathematical derivation of the Weibull distribution. This was inspired by the broken stick model widely used in ecology in studying species-abundance curves.

Fargo et al. (1982) used the PDF and CDF of the Weibull distribution to fit the within-tree population of the studied tree species. Hogg and Nordheim (1983) used modified Weibull distribution to obtain survival curves of the eggs and larvae of the studied species while Luo et al. (2015) and Baek et al. (2017) used Weibull distribution to obtain the survival curve of their studied species. Legge and Krupa (1989) used the Weibull family of distribution to model air pollution data. Newman and Aplin (1992) observed that the Weibull distribution is the best in fitting survival time modeling in toxicology.

Nanos and Montero (2002), Sarkkola et al. (2003), Neeff et al. (2003), Sarkkola et al. (2005), Wang et al. (2010), Chan et al. (2011), Forrester et al. (2013) and Navarro-Cerrillo et al. (2014) used the Weibull distribution to model and predict the diameter distributions of stand. Tjørve (2003) noted from review that the Weibull distribution is a sigmoid model that can prove useful in modeling species-area relationships. He et al. (2004) used Weibull distribution and five other probability distributions to characterized patch size distributions of their studied district. In that research, lognormal distribution performed better in fitting the data than the other five distributions. MacKenzie et al. (2004) applied the Weibull distribution to model the densities and diversity of invertebrates at Lake Michigan and the abiotic factors that affect the dynamics of the invertebrates. Yang et al. (2004) proposed a model with Weibull and exponential models as special case, which can be applied in the modeling of fire disturbances in forest landscapes. The model was an improvement over the exponential and Weibull fire interval distribution as it accounts for the separation of fire ignition from fire occurrence. Eliopoulos et al. (2005) used the Weibull distribution in the estimation of the survival rate of the studied parasitoid when subjected to temperature and other factors. Al-Saidy et al. (2005) used two forms of Weibull distributions to estimate the bias of three measures of overlap used in quantitative ecology. McKenzie et al. (2006) used Weibull and other distributions to model and characterize some factors that affect the spatial and temporal distribution of fire occurrence in forests. In that study, Weibull distribution was used to fit real and simulated exhibited different parameters in modeling temporal trends in fire hazards. Carlson et al. (2008) used the three parameter Weibull distribution to determine the relationship between the changes in diameter distribution and mid-rotation fertilization. Kapur et al. (2008) developed dynamic stock and flow model, utilizing the gamma, lognormal and Weibull distributions to estimate the in-use cement stocks in the United States. Moritz et al. (2009) observed using the cedar fire in 2003 in southern California as a case study, that the parameters estimates of the Weibull distribution was able to fit adequately the previous fire frequency analysis of the shrubland even with the use of censored data.

Bonou et al. (2009) used the Weibull distribution to model the stem diameter and height structure of the endangered tree species found in the four communities in the Lama Forest reserve of Benin Republic in West Africa. Li et al. (2011) observed that the Weibull and gamma distributions performed better than the normal distribution in simulating the distribution of leaf and branch. Huffman et al. (2012) used Weibull distribution to model the interrelationship among key structural elements and time since fire in forest ecological management. Studds et al. (2012) concluded from his research that Weibull distribution provides best fits for assessing the probability of dispersal across the breeding range of their studied bird species.

Parameter estimates from strongly skewed probability distributions are often encountered in ecology, for example in diameter stands of trees. Most often ecologists are confronted with the tasks of interpreting ecological results. This is due to the variability of the underlying parameters of the distributions that was used to fit the data. This is further complicated by the confounding effects of the parameters due to errors in measurement and experimentation. In order to minimize the effects of the aforementioned problems, Taubert et al. (2013) investigated the bias nature of the parameters of three frequently used probability distributions namely; Weibull, negative exponential and power-law distributions.

Akinci et al. (2013), Guseinoviene et al. (2014) and Sedaghat et al. (2016) used the Weibull distribution to estimate and model wind speed. Jankowski et al. (2013) used Weibull distribution to determine viral count in a disease ecological study. Zabel et al. (2014) used modified Weibull, Weibull and exponential distributions to model migrating adult salmonids. In a study of different growth curve models, Dasgupta (2015) noted that growth curves of harvesting and forecast of market supply of yam can be approximated by the Weibull distribution. Gao and Perry (2016) used the cumulative Weibull regression to model the bio-geographical patterns of their study subjects. Subedi and Fox (2016) used Weibull distribution to model soil fertility rating used to predict the effects of fertilizer to the growth of loblolly pine.

In a recent study, Rogeau and Armstrong (2017) noted that the use of Weibull distribution in modeling the effect of topography on wildfire distribution in many studies might have left out some effects unexplored. All these contributions indicate the fact that Weibull distribution has been applied in ecological studies.

1.2 Ordinary Differential Equations of Probability Functions

Differential equations often arise from the understanding and modeling of real life problems or some observed physical phenomena. Newton and Leibniz are believed to be the inventors, obtained from calculus. Types of differential equations include: ordinary differential equations, partial differential equations, stochastic differential equations, fractional differential equations, neural differential equations, delay differential equations and so on.

In science, engineering and social sciences, mathematical models are developed for the understanding of physical phenomena. In view of that Enszer et al. (2011) stated that modeling dynamic systems is challenging because of the inherent difficulty of incorporating all the uncertainties in the parameters into mathematical models.

However, when the physical phenomena are characterized by uncertainties, then statistical models are used. This is because some phenomena often follow the same pattern and orientation; they are therefore grouped into different general models called probability density functions. The observed events may be discrete or continuous. Bernoulli, Geometric, binomial, negative binomial, hypergeometric, Rademacher, discrete uniform and Poisson distributions are few examples of the discrete case while the normal, reciprocal, trapezoidal, Nakagami, Rice, Pareto, Gumbel, exponential, Johnson SU, gamma, Chi-square, logit-normal and beta distributions are few examples of the continuous case. In comparing deterministic and probabilistic models, Calderhead and Girolami (2011)

emphasized on the usefulness of probabilistic models, such as uncertainty quantification, predictions and hypothesis tests on the parameters of the models.

Ordinarily, derivatives are used in the estimation of parameters of the probability density functions. The method of maximum likelihood is an example (Akaike, 1998).

Ordinary differential equations (ODE) as an extension to derivatives are indispensable in the understanding of physical, social and biological processes. Most often the ODE does not have analytic solutions and researchers resort to numerical methods for approximate solutions. One of the reasons why the analytic solutions do not always exist is the presence of uncertainties. Over the centuries, researchers have classified and grouped these uncertainties into probability density functions. In other to fully quantify uncertainties, other functions were created, such as the cumulative distribution function, Quantile function, survival function, inverse survival function, odd function, hazard function and reversed hazard function. Incorporation of uncertainties in mathematical modeling has been useful in ecology, engineering, epidemiology, disease control, demography, psephology, meteorology, astronomy, medicine, geology, finance, ecology, biogeography, sociology, economics and so on.

ODE has been helpful in statistical and probabilistic models. It can also be seen as a tool for measuring uncertainties and prediction for example in engineering analysis (Papadimitriou, Katafygiotis and Beck (1995). The outcome is that the solution of the ODE if it exists is the probability density function of the model considered. This was the outcome of the result obtained by (Steinsaltz et al., 2005), when in modeling mutation as it applied to aging, furthermore, the solution of the ODE they formulated for their model converged to the probability density function of the distribution that they considered.

Interestingly, the convergence of solutions are not restricted to the probability density function only, it can be extended to the following: kernel density of the distribution (Shotorban, 2000), the parameters of the systems defined by the ODEs (Atencia and Joya, 2011) and probability generating function (Reed and Hughes, 2002).

However, there seems to be an argument on the estimation by ODE and the maximum likelihood, the details can be found in Hirose (2011, 2012). The maximum likelihood is limited when the parameters that defined the model are huge and is safer to estimate the parameters effectively by the use of ODE. The existence and uniqueness of solutions of the ODEs of the models have been considered (Knopoff, 2013), however, Ghitany et al. (2011) restricted their research scope to the maximum likelihood estimators MLEs of the parameters of the class of distribution that they considered.

The limitation of the use of ODE in modeling uncertainties is that it does not take into account, the time factor (Hall and Gandar, 1996), and the inability of the uncertainties that characterizes a model to be fully modeled using the ODEs. This was also a remark made by Banks et al. (2003), as they developed some methods of incorporating uncertainties and variability into systems that cannot be reduced to ODEs. It is more expensive to incorporate uncertainties into models than to simply fit the data into well-known probability distributions.

The aim of this research is to develop homogenous ordinary differential equations for the probability density function (PDF), Quantile function (QF), survival function (SF), inverse survival function (ISF),

hazard function (HF) and reversed hazard function (RHF) of Weibull distribution as it is observed to be commonly applied in ecological research. This will also help to provide the answers as to whether there are discrepancies between the support of the Weibull distribution and the necessary conditions for the existence of the ODEs.

1.3 Weibull Distribution

This is a continuous and lifetime distribution widely used in survival and lifetime analysis. The distribution was named after Swedish mathematician; Ernst Hjalmar Waloddi Weibull. The various aspects of the Weibull distribution has been studied or reported in literature. These include: maximum likelihood estimation (Menon, 1963; Lemon, 1975); Cohen, 1965), inferences on the parameters (Thoman et al., 1969), qualitative reviews (Hallinan, 1993; Pham and Lai, 2007), moments and moments of order statistics (Lieblein, 1955), statistical tests (Littell et al., 1979).

The distribution has witnessed generations, modifications, compounding, extensions and parameterization such as: discrete Weibull distribution by Nakagawa and Osaki (1975), generalization by Mudholkar et al. (1996), exponentiated Weibull distribution by Nassar and Eissa (2003), Nadarajah et al. (2013) and Shittu and Adepoju (2014), Normalized Weibull distribution by Marabi et al. (2003), Beta-Weibull distribution by Lee et al. (2007). Weibull-geometric distribution by Barreto-Souza et al. (2011), Transmuted Weibull distribution by Aryal and Tsokos (2011), beta generalized Weibull distribution by Singla et al. (2012), beta Weibull Poisson distribution by Percontini et al. (2013), Mixture Weibull distributions by Razali and Al-Wakeel (2013), exponentiated generalized Weibull distribution by Adepoju et al. (2013).

Also available are: Kumaraswamy modified Weibull distribution by Cordeiro et al. (2014), Weibull-exponential distribution by Oguntunde et al. (2015), transmuted exponential-Weibull distribution by Saboor et al. (2015), Marshall-Olkin exponential Weibull distribution by Pogány et al. (2015), Weibull-Lomax distribution by Tahir et al. (2015), Weighted weibull distribution by Dey et al. (2015), additive Weibull-Geometric Distribution by Elbatal et al. (2016), Weibull power function distribution by Tahir et al. (2016 a), Weibull-Pareto distribution by Tahir et al (2016 b), Weibull-Dagum distribution by Tahir et al. (2016 c), Gamma Dual Weibull Model by Castellares and Lemonte (2016), Marshall-Olkin gamma-Weibull distribution by Saboor and Pogany (2016), new Weibull-G family of distributions by Tahir et al. (2016 d), Kumaraswamy generalized power Weibull distribution by Selim and Badr (2016).

It is a widely applied probability distribution such as: modeling the chances of occurrence of earthquake by Hagiwara (1974), analysis of survival data in ecology by Pinder et al. (1978), inventory control by Tadikamalla (1978), forecasting of change in technology by Sharif and Islam (1980), ferrography by Roylance and Pocock (1983), metamorphosis of insects by Wagner et al. (1984), hydrology by Singh (1987), estimation of wind speed by Seguro and Lambert (2000), modeling preservation methods by Mafart et al. (2002) and Bai et al. (2013), modeling cracks in rocks by Wong et al. (2006), assessing wind power density by Mohammadi et al. (2016).

2. MATERIAL AND METHODS

Ordinary differential calculus was used to obtain the ODEs whose solutions are the PDF, QF, SF, ISF, HF and RHF of the Weibull distributions respectively.

3. RESULTS

This section contains the detailed results.

3.1 Probability Density Function

The probability density function of the Weibull distribution is given as;

$$f(x) = \frac{\eta}{\sigma} \left(\frac{x}{\sigma}\right)^{\eta-1} e^{-\left(\frac{x}{\sigma}\right)^\eta} \quad (1)$$

The PDF is characterized by the shape parameter $\sigma > 0$, scale parameter $\eta > 0$ and the support $x > 0$.

To obtain the first order ordinary differential equation for the probability density function of the Weibull distribution, differentiate equation (1), to obtain;

$$f'(x) = \left\{ \frac{\frac{\eta-1}{\sigma} \left(\frac{x}{\sigma}\right)^{\eta-2} - \frac{\eta}{\sigma} \left(\frac{x}{\sigma}\right)^{\eta-1} e^{-\left(\frac{x}{\sigma}\right)^\eta}}{\left(\frac{x}{\sigma}\right)^{\eta-1} e^{-\left(\frac{x}{\sigma}\right)^\eta}} \right\} f(x) \quad (2)$$

$$f'(x) = \left\{ \frac{\eta-1}{x} - \frac{\eta}{\sigma} \left(\frac{x}{\sigma}\right)^{\eta-1} \right\} f(x) \quad (3)$$

The condition necessary for the existence of the equation is $x, \eta, \sigma > 0$.

The differential equations can only be obtained for particular values of η and σ .

When $\eta = 1$, equation (3) becomes;

$$f'_a(x) = \left(-\frac{1}{\sigma}\right) f_a(x) \quad (4)$$

$$\sigma f'_a(x) + f_a(x) = 0 \quad (5)$$

When $\eta = 2$, equation (3) becomes;

$$f'_b(x) = \left\{ \frac{1}{x} - \frac{2}{\sigma} \left(\frac{x}{\sigma}\right) \right\} f_b(x) \quad (6)$$

$$\sigma^2 x f'_b(x) + (2x^2 - \sigma^2) f_b(x) = 0 \quad (7)$$

When $\eta = 3$, equation (3) becomes;

$$f'_c(x) = \left\{ \frac{2}{x} - \frac{3}{\sigma} \left(\frac{x}{\sigma}\right)^2 \right\} f_c(x) \quad (8)$$

$$\sigma^3 x f'_c(x) + (3x^3 - 2\sigma^3) f_c(x) = 0 \quad (9)$$

To obtain a simplified ordinary differential equation, differentiate equation (3);

$$f''(x) = \left(\frac{\eta-1}{x} - \frac{\eta}{\sigma} \left(\frac{x}{\sigma} \right)^{\eta-1} \right) f'(x) - \left(\frac{(\eta-1)}{x^2} + \frac{\eta}{\sigma} \left(\frac{\eta-1}{\sigma} \right) \left(\frac{x}{\sigma} \right)^{\eta-2} \right) f(x) \quad (10)$$

The condition necessary for the existence of the equation is $x, \eta, \sigma > 0$.

The following equations obtained from (3) are needed to simplify equation (10);

$$\frac{f'(x)}{f(x)} = \frac{\eta-1}{x} - \frac{\eta}{\sigma} \left(\frac{x}{\sigma} \right)^{\eta-1} \quad (11)$$

$$\frac{\eta}{\sigma} \left(\frac{x}{\sigma} \right)^{\eta-1} = \frac{\eta-1}{x} - \frac{f'(x)}{f(x)} \quad (12)$$

$$\frac{\eta}{\sigma} \left(\frac{\eta-1}{\sigma} \right) \left(\frac{x}{\sigma} \right)^{\eta-2} = \left(\frac{\eta-1}{x} \right) \left(\frac{\eta-1}{x} - \frac{f'(x)}{f(x)} \right) \quad (13)$$

Substitute equations (11) and (13) into equation (10) to obtain;

$$f''(x) = \frac{f'^2(x)}{f(x)} - \left(\frac{(\eta-1)}{x^2} + \left(\frac{\eta-1}{x} \right) \left(\frac{\eta-1}{x} - \frac{f'(x)}{f(x)} \right) \right) f(x) \quad (14)$$

$$f''(x) = \frac{f'^2(x)}{f(x)} - \frac{(\eta-1)f(x)}{x^2} - \frac{(\eta-1)^2 f(x)}{x^2} + \frac{(\eta-1)f'(x)}{x} \quad (15)$$

The second order differential equation for the probability density function of the Weibull distribution is given by;

$$x^2 f(x) f''(x) - x^2 f'^2(x) + n(n-1) f^2(x) - (n-1) x f(x) f'(x) = 0 \quad (16)$$

$$f(1) = \frac{\eta}{\sigma^\eta} e^{-\left(\frac{1}{\sigma}\right)^\eta} \quad (17)$$

$$f'(1) = \frac{\eta(\eta-1)\sigma^2 - \eta}{\sigma^{2\eta}} e^{-\left(\frac{1}{\sigma}\right)^\eta} \quad (18)$$

3.2 Quantile Function

The Quantile function of the Weibull distribution is given as;

$$Q(p) = \sigma (-\ln(1-p))^\eta \quad (19)$$

The QF is characterized by the shape parameter $\sigma > 0$, scale parameter $\eta > 0$ and the support $0 < p < 1$.

Differentiate equation (19);

$$Q'(p) = \frac{\sigma \eta}{1-p} (-\ln(1-p))^{\eta-1} \quad (20)$$

The condition necessary for the existence of the equation is $\eta, \sigma > 0, 0 < p < 1$.

Substitute equation (19) into equation (20);

$$Q'(p) = \frac{\eta}{1-p} \frac{Q(p)}{(-\ln(1-p))} \quad (21)$$

Simplify equation (19) to obtain;

$$\left(\frac{Q(p)}{\sigma} \right)^{\frac{1}{\eta}} = -\ln(1-p) \quad (22)$$

Substitute equation (22) into (21);

$$Q'(p) = \frac{\eta \sigma^{\frac{1}{\eta}}}{1-p} Q^{\frac{\eta-1}{\eta}}(p) \quad (23)$$

The first order differential equation for the Quantile function of the Weibull distribution is given by

$$(1-p)Q'(p) - \eta \sigma^{\frac{1}{\eta}} Q^{\frac{\eta-1}{\eta}}(p) = 0 \quad (24)$$

$$Q(0) = 0 \quad (25)$$

Some special cases of equation (24) is considered;

When $\eta = 1$, equation (24) becomes;

$$(1-p)Q'_a(p) = 0 \quad (26)$$

When $\eta = 2$, equation (24) becomes;

$$(1-p)Q'_b(p) - 2\sigma^{\frac{1}{2}} Q^{\frac{1}{2}}(p) = 0 \quad (27)$$

3.3 Survival Function

The survival function of the Weibull distribution is given as;

$$S(t) = e^{-\left(\frac{t}{\sigma}\right)^{\eta}} \quad (28)$$

The SF is characterized by the shape parameter $\sigma > 0$, scale parameter $\eta > 0$ and the support $x > 0$.

Differentiate equation (28);

$$S'(t) = -\frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} e^{-\left(\frac{t}{\sigma}\right)^{\eta}} = -f(t) \quad (29)$$

The condition necessary for the existence of the equation is $t, \eta, \sigma > 0$.

$$S'(t) + f(t) = 0 \quad (30)$$

$$S(1) = e^{-\left(\frac{1}{\sigma}\right)^{\eta}} \quad (31)$$

Also, equation (29) can be simplify using equation (28).

$$S'(t) = -\frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} S(t) \quad (32)$$

The differential equations can only be obtained for particular values of η and σ .

When $\eta = 1$, equation (32) becomes;

$$S'_a(t) = -\frac{1}{\sigma} S_a(t) \quad (33)$$

$$\sigma S'_a(t) + S_a(t) = 0 \quad (34)$$

When $\eta = 2$, equation (32) becomes;

$$S'_b(t) = -\frac{2}{\sigma} \left(\frac{t}{\sigma}\right) S_b(t) \quad (35)$$

$$\sigma^2 S'_b(t) + 2t S_b(t) = 0 \quad (36)$$

When $\eta = 3$, equation (32) becomes;

$$S'_c(t) = -\frac{3}{\sigma} \left(\frac{t}{\sigma}\right)^2 S_c(t) \quad (37)$$

$$\sigma^3 S'_c(t) + 3t^2 S_c(t) = 0 \quad (38)$$

Using the results of the probability density function to obtain the one for the survival function;
Modifying equation (3);

$$S''(t) = \left\{ \frac{\eta-1}{t} - \frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} \right\} S'(t) \quad (39)$$

The differential equations can only be obtained for particular values of η and σ .

When $\eta = 1$, equation (39) becomes;

$$S''_d(t) = \left(-\frac{1}{\sigma}\right) S'_d(t) \quad (40)$$

$$\sigma S''_d(t) + S'_d(t) = 0 \quad (41)$$

When $\eta = 2$, equation (39) becomes;

$$S''_e(t) = \left\{ \frac{1}{t} - \frac{2}{\sigma} \left(\frac{t}{\sigma}\right) \right\} S'_e(t) \quad (42)$$

$$\sigma^2 t S''_e(t) + (2t^2 - \sigma^2) S'_e(t) = 0 \quad (43)$$

When $\eta = 3$, equation (39) becomes;

$$S''_f(t) = \left\{ \frac{2}{t} - \frac{3}{\sigma} \left(\frac{t}{\sigma}\right)^2 \right\} S'_f(t) \quad (44)$$

$$\sigma^3 t S''_f(t) + (3t^3 - 2\sigma^3) S'_f(t) = 0 \quad (45)$$

Alternatively, using equations (31) and (32) on (39) yields a different ordinary differential equation

$$S''(t) = \left\{ \frac{\eta-1}{t} - \frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} \right\} \times -\frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} \times S(t) \quad (46)$$

$$S''(t) = -\frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} \left\{ \frac{\eta-1}{t} - \frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} \right\} S(t) \quad (47)$$

The differential equations can only be obtained for particular values of η and σ .

When $\eta = 1$, equation (47) becomes;

$$S_g''(t) = -\frac{1}{\sigma} \left\{ -\frac{1}{\sigma} \right\} S_g(t) \quad (48)$$

$$\sigma^2 S_g''(t) - S_g(t) = 0 \quad (49)$$

When $\eta = 2$, equation (47) becomes;

$$S_h''(t) = -\frac{2}{\sigma} \left(\frac{t}{\sigma} \right) \left\{ \frac{1}{t} - \frac{2}{\sigma} \left(\frac{t}{\sigma} \right) \right\} S_h(t) \quad (50)$$

$$\sigma^4 S_h''(t) - 2(2t^2 - \sigma^2) S_h(t) = 0 \quad (51)$$

When $\eta = 3$, equation (47) becomes;

$$S_i''(t) = -\frac{3}{\sigma} \left(\frac{t}{\sigma} \right)^2 \left\{ \frac{2}{t} - \frac{3}{\sigma} \left(\frac{t}{\sigma} \right)^2 \right\} S_i(t) \quad (52)$$

$$\sigma^6 S_i''(t) - 3t(3t^3 - 2\sigma^3) S_i(t) = 0 \quad (53)$$

Furthermore, the third order ordinary differential equations can be obtained by the modification of equation (16);

$$t^2 S'(t) S'''(t) - t^2 S''^2(t) + n(n-1) S'^2(t) - (n-1) t S'(t) S''(t) = 0 \quad (54)$$

$$S'(1) = -\frac{\eta}{\sigma^\eta} e^{-\left(\frac{1}{\sigma}\right)^\eta} \quad (55)$$

$$S''(1) = -\frac{\eta(\eta-1)\sigma^2 - \eta}{\sigma^{2\eta}} e^{-\left(\frac{1}{\sigma}\right)^\eta} \quad (56)$$

3.4 Inverse Survival Function

The inverse survival function of the Weibull distribution is given as;

$$Q(p) = \sigma \left(\ln \left(\frac{1}{p} \right) \right)^{\frac{1}{\eta}} \quad (57)$$

The ISF is characterized by the shape parameter $\sigma > 0$, scale parameter $\eta > 0$ and the support $0 < p < 1$.

Differentiate equation (57);

$$Q'(p) = -\frac{\sigma}{\eta p} \left(\ln \left(\frac{1}{p} \right) \right)^{\frac{1}{\eta} - 1} \quad (58)$$

$$Q'(p) = -\frac{\sigma \left(\ln \left(\frac{1}{p} \right) \right)^{\frac{1}{\eta}}}{\eta p \left(\ln \left(\frac{1}{p} \right) \right)} \quad (59)$$

The condition necessary for the existence of the equation is $\eta, \sigma > 0, 0 < p < 1$.

Substitute equation (57) into equation (59);

$$Q'(p) = -\frac{Q(p)}{\eta p \left(\ln \left(\frac{1}{p} \right) \right)} \quad (60)$$

Simplify equation (57) to obtain;

$$\left(\frac{Q(p)}{\sigma}\right)^\eta = \ln\left(\frac{1}{p}\right) \quad (61)$$

Substitute equation (61) into (60);

$$Q'(p) = -\frac{\sigma^\eta Q^{1-\eta}(p)}{\eta p} \quad (62)$$

$$\eta p Q'(p) + \sigma^\eta Q^{1-\eta}(p) = 0 \quad (63)$$

The ordinary differential equations can be obtained for particular values of the parameters of equation (63). Some cases are considered.

Table 1: First order ODEs of the ISF of Weibull Distribution for different parameters.

η	σ	Ordinary Differential Equation
1	1	$pQ'(p) + 1 = 0$
1	2	$pQ'(p) + 2 = 0$
2	1	$2pQ(p)Q'(p) + 1 = 0$
2	2	$pQ(p)Q'(p) + 2 = 0$

3.5 Hazard Function

The hazard function of the Weibull distribution is given as;

$$h(t) = \frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} \quad (64)$$

The HF is characterized by the shape parameter $\sigma > 0$, scale parameter $\eta > 0$ and the support $x > 0$.

Differentiate equation (64);

$$h'(t) = \frac{\eta}{\sigma} \left(\frac{\eta-1}{\sigma}\right) \left(\frac{t}{\sigma}\right)^{\eta-2} = \frac{\eta(\eta-1)}{\sigma^2} \left(\frac{t}{\sigma}\right)^{\eta-2} \quad (65)$$

The condition necessary for the existence of the equation is $t, \eta, \sigma > 0$.

Substitute equation (64) into equation (65);

$$h'(t) = \frac{\eta-1}{t} h(t) \quad (66)$$

The first order ordinary differential equation for the Hazard function of the Weibull distribution is given as;

$$th'(t) - (\eta-1)h(t) = 0 \quad (67)$$

$$h(1) = \frac{\eta}{\sigma^\eta} \quad (68)$$

Differentiate equation (66);

$$h''(t) = \frac{\eta(\eta-1)(\eta-2)}{\sigma^3} \left(\frac{t}{\sigma}\right)^{\eta-3} \quad (69)$$

The condition necessary for the existence of the equation is $t, \eta, \sigma > 0$.

Two ordinary differential equations can be obtained from the simplification of equation (69),

ODE 1; Use equation (64) in (69);

$$h''(t) = \frac{(\eta-1)(\eta-2)}{t^2} h(t) \quad (70)$$

$$t^2 h''(t) - (\eta-1)(\eta-2)h(t) = 0 \quad (71)$$

ODE 2; Use equation (65) in (69);

$$h''(t) = \frac{(\eta-2)}{t} h'(t) \quad (72)$$

$$t h''(t) - (\eta-2)h'(t) = 0 \quad (73)$$

$$h'(1) = \frac{\eta(\eta-1)}{\sigma^\eta} \quad (74)$$

Differentiate equation (69);

$$h'''(t) = \frac{\eta(\eta-1)(\eta-2)(\eta-3)}{\sigma^4} \left(\frac{t}{\sigma}\right)^{\eta-4} \quad (75)$$

The condition necessary for the existence of the equation is $t, \eta, \sigma > 0$.

Three ordinary differential equations can be obtained from the simplification of equation (75),

ODE 1; Use equation (64) in (75);

$$h'''(t) = \frac{(\eta-1)(\eta-2)(\eta-3)}{t^3} h(t) \quad (76)$$

$$t^3 h'''(t) - (\eta-1)(\eta-2)(\eta-3)h(t) = 0 \quad (77)$$

ODE 2; Use equation (65) in (75);

$$h'''(t) = \frac{(\eta-2)(\eta-3)}{t^2} h'(t) \quad (78)$$

$$t^2 h'''(t) - (\eta-2)(\eta-3)h'(t) = 0 \quad (79)$$

ODE 3; Use equation (69) in (75);

$$h'''(t) = \frac{(\eta-3)}{t} h''(t) \quad (80)$$

$$t h'''(t) - (\eta-3)h''(t) = 0 \quad (81)$$

$$h''(1) = \frac{\eta(\eta-1)(\eta-2)}{\sigma^\eta} \quad (82)$$

3.6 Reversed Hazard Function

The reversed hazard function of the Weibull distribution is given as;

$$j(t) = \frac{\frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} e^{-\left(\frac{t}{\sigma}\right)^\eta}}{1 - e^{-\left(\frac{t}{\sigma}\right)^\eta}} \quad (83)$$

The RHF is characterized by the shape parameter $\sigma > 0$, scale parameter $\eta > 0$ and the support $x > 0$.

To obtain the first order ordinary differential equation for the reversed hazard function of the Weibull

distribution, differentiate equation (83), to obtain;

$$j'(t) = \left\{ \frac{\frac{\eta-1}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-2} - \frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} e^{-\left(\frac{t}{\sigma}\right)^\eta} - \frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} e^{-\left(\frac{t}{\sigma}\right)^\eta} (1 - e^{-\left(\frac{t}{\sigma}\right)^\eta})^{-2}}{\left(\frac{t}{\sigma}\right)^{\eta-1} e^{-\left(\frac{t}{\sigma}\right)^\eta} (1 - e^{-\left(\frac{t}{\sigma}\right)^\eta})^{-1}} \right\} j(t) \quad (84)$$

$$j'(t) = \left\{ \frac{\eta-1}{t} - \frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} - \frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} e^{-\left(\frac{t}{\sigma}\right)^\eta}}{1 - e^{-\left(\frac{t}{\sigma}\right)^\eta}} \right\} j(t) \quad (85)$$

The condition necessary for the existence of the equation is $t, \eta, \sigma > 0$.

$$j'(t) = \left(\frac{\eta-1}{t} - \frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} - j(t) \right) j(t) \quad (86)$$

The differential equations can only be obtained for particular values of η and σ . Some cases are given in Table 2.

Table 2: First order ODEs of the ISF of Weibull Distribution for different parameters.

η	σ	Ordinary Differential Equation
1	1	$j'(t) + j(t) + j^2(t) = 0$
1	2	$2j'(t) + j(t) + 2j^2(t) = 0$
2	1	$tj'(t) + (2t^2 - 1)j(t) + tj^2(t) = 0$
2	2	$2tj'(t) + (t^2 - 2)j(t) + 2tj^2(t) = 0$

To obtain the second order ordinary differential equation, differentiate equation (86);

$$j''(t) = \left(\frac{\eta-1}{t} - \frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} - j(t) \right) j'(t) - \left(\frac{(\eta-1)}{t^2} + \frac{\eta}{\sigma} \left(\frac{\eta-1}{\sigma}\right) \left(\frac{t}{\sigma}\right)^{\eta-2} + j'(t) \right) j(t) \quad (87)$$

The following equations obtained from (86) are needed to simplify equation (87);

$$\frac{j'(t)}{j(t)} = \frac{\eta-1}{t} - \frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} - j(t) \quad (88)$$

$$\frac{\eta}{\sigma} \left(\frac{t}{\sigma}\right)^{\eta-1} = \frac{\eta-1}{t} - \frac{j'(t)}{j(t)} - j(t) \quad (89)$$

$$\frac{\eta}{\sigma} \left(\frac{\eta-1}{\sigma}\right) \left(\frac{t}{\sigma}\right)^{\eta-2} = \left(\frac{\eta-1}{t}\right) \left(\frac{\eta-1}{t} - \frac{j'(t)}{j(t)} - j(t)\right) \quad (90)$$

Substitute equations (88) and (90) into equation (87) to obtain;

$$j''(t) = \frac{j'^2(t)}{j(t)} - \left(\frac{(\eta-1)}{t^2} + \left(\frac{\eta-1}{t}\right) \left(\frac{\eta-1}{t} - \frac{j'(t)}{j(t)} - j(t)\right) + j'(t) \right) j(t) \quad (91)$$

The differential equations can only be obtained for particular values of η and σ .

3.7 Application

Weibull distribution was used to fit the daily radiation data of Port Harcourt Nigeria. The data was obtained from Nigeria Meteorological Agency. It is for the month of September 2015. The data is presented;

Table 3: Radiation Data of Port Harcourt Nigeria

Date	Radiation level	Date	Radiation level	Date	Radiation level
9/1/2015	109.17	9/11/2015	164.41	9/21/2015	178.88
9/2/2015	77.60	9/12/2015	127.58	9/22/2015	184.14
9/3/2015	132.84	9/13/2015	86.81	9/23/2015	176.25
9/4/2015	109.17	9/14/2015	118.38	9/24/2015	103.91
9/5/2015	52.61	9/15/2015	114.43	9/25/2015	80.23
9/6/2015	73.66	9/16/2015	90.75	9/26/2015	203.87
9/7/2015	149.94	9/17/2015	81.55	9/27/2015	194.66
9/8/2015	147.31	9/18/2015	119.69	9/28/2015	143.37
9/9/2015	110.48	9/19/2015	170.99	9/29/2015	107.85
9/10/2015	143.37	9/20/2015	155.20	9/30/2015	140.74

The descriptive statistics of the data are summarized in Table 4.

Table 4: Descriptive Statistics of the Radiation Data

Mean	median	Standard Deviation	Variance	Skewness	Kurtosis	Range	Minimum	Maximum
128.33	123.64	39.28	1542.76	0.117	-0.772	151.26	52.61	203.87

The data was fitted with the Weibull and the estimated values of the parameters are given in Table 5.

Table 5: Parameter Estimates of the Data Fit

Parameter	Estimated Value	Standard Deviation
Shape	3.707377054	0.531378272
Scale	142.4835373	7.408452151

The estimates are substituted in equations (3), (24), (33), (63), (67) and (86). The solutions are the PDF, QF, SF, ISF, HF and RHF of the data. For example, the case of hazard function was considered.

Substitute $\eta = 3.7$ and $\sigma = 142.5$ into equations (67) and (68) to obtain;

$$th'(t) - 2.7h(t) = 0 \quad (92)$$

$$h(1) = \frac{3.7}{142.5^{3.7}} = 0.00000004 \quad (93)$$

The solution gives the hazard function of the Weibull distribution.

4. CONCLUSION

The ODEs of the probability functions whose solutions are the respective probability functions of the Weibull distribution have been obtained. This result has shown that the ODE is not limited to the PDF but other probability functions. The usefulness of the ODEs in fitting is still vague and need further exploration. The research is crucial in ecological studies as the study of nature of the studied distribution can be extended to the behavior of the ODEs that define them. The parameters determine not only the distribution but also the nature of the ODE. Finally the results are similar to the series obtained by Okagbue et al., (2017 a, b, c, d, e, f, g, h, i, j, k, l, m, n, o). In addition some methods of solutions to the various ODE may be explored such as Anake et al, (2012 a, b, c, 2013, 2015).

5. REFERENCES

Adepoju, K. A., Shittu, O. I., Tijani I. A., 2013, The Statistical Properties of the Exponentiated Generalized Weibull Distribution with Applications. *Proc. Annual the Confe. of the Nig. Stat.* 136- 143.

Akaike, H., 1998, Information theory and an extension of the maximum likelihood principle. In *Selected Papers of Hirotugu Akaike* (pp. 199-213). Springer New York.

Akinci, T.C., Seker, S., Guseinoviene, E., Nayir, A., 2013, Statistical analysis and Hurst parameter estimation for wind speed in Kırklareli area of Turkey. *8th Int. Conf. Exhibition on Ecological Vehicles and Renewable Energies, EVER*. Article number 6521592.

Al-Saidy, O., Samawi, H.M., Al-Saleh, M.F., 2005, Inference on overlap coefficients under the Weibull distribution: Equal shape parameter. *ESAIM – Prob. Stat.* **9**, 206-219.

Anake, T. A., Awoyemi, D. O., Adesanya, A. A., 2012, A one step method for the solution of general second order ordinary differential equations. *International Journal of science and Technology*, 2(4), 159-163.

Anake, T. A., Awoyemi, D. O., Adesanya, A. O., 2012, One-step implicit hybrid block method for the direct solution of general second order ordinary differential equations. *IAENG International Journal of Applied Mathematics*, 42(4), 224-228.

Anake, T. A., Awoyemi, D. O., Adesanya, A. A., Famewo, M. M., 2012, Solving general second order ordinary differential equations by a one-step hybrid collocation method. *International Journal of Science and Technology*, 2(4), 164-168.

Anake, T. A., Adesanya, A. O., Oghonyon, J. G., Agarana, M. C., 2013, Block algorithm for general third order ordinary differential equation. *Icastor Journal of Mathematical Sciences*, 7(2), 127-136.

- Anake, T. A., Bishop, S. A., Agboola, O. O., 2015, On a hybrid numerical algorithm for the solutions of higher order ordinary differential equations. *TWMS J. Pure Appl. Math.*, 6(2).
- Aryal, G. R., Tsokos, C. P., 2011, Transmuted Weibull distribution: A generalization of the Weibull probability distribution. *Euro. J. Pure Appl. Math*, 4, 89-102.
- Atencia, M., Joya, G., 2011, Statistical properties of the 'Hopfield estimator' of dynamical systems. *ESANN 2011, 19th European Symposium on Artificial Neural Networks, Computational Intelligence and Machine Learning*, 47-52.
- Baek, S., Hwang, A., Kim, H., Lee, H., Lee, J.-H., 2017, Temperature-dependent development and oviposition models of *Halyomorpha halys* (Hemiptera: Pentatomidae). *J. Asia-Pacific Entomology*. **20**, 367-375.
- Bai, J., Wang, J., Xiao, H., Ju, H., Liu, Y., Gao, Z., 2013, Weibull distribution for modeling drying of grapes and its application. *Transac. Chinese Soc. Agric. Engine*. **29**, 278-285.
- Balakrishnan, N., Lai, C. D., 2009, Continuous Bivariate Distributions, 2nd edition, Springer New York, London.
- Banks, H. T., Bortz, D. M., Holte, S. E., 2003, Incorporation of variability into the modeling of viral delays in HIV infection dynamics. *Math. Biosciences*. **183**, 63-91.
- Barreto-Souza, W., de Morais, A. L., Cordeiro, G. M., 2011, The Weibull-geometric distribution. *J. Stat. Comput. Simul.* **81**, 645-657.
- Bonou, W., Glèlè Kakaï, R., Assogbadjo, A.E., Fonton, H.N., Sinsin, B., 2009, Characterisation of *Afzelia africana* Sm. habitat in the Lama forest reserve of Benin. *Forest Ecol. Magt.* **258**, 1084-1092.
- Calderhead, B., Girolami, M., 2011, Statistical analysis of nonlinear dynamical systems using differential geometric sampling techniques. *Interface Focus*. **1**, 821-835.
- Carlson, C.A., Burkhart, H.E., Allen, H.L., Fox, T.R., 2008, Absolute and relative changes in tree growth rates and changes to the stand diameter distribution of *Pinus taeda* as a result of midrotation fertilizer applications. *Canadian J. Forest Res.* **38**, 2063-2071.
- Castellares, F., Lemonte, A. J., 2016, On the Gamma Dual Weibull Model. *Amer. J. Math. Magt. Sci.* **35**, 124-132.
- Chan, M.-H., Chen, J.-H., Lin, H.C., Fujimoto, N., 2011, Tree ring analysis for growth pattern and age structure of the subtropical broad-leaved forest in Taiwan. *J. Faculty Agric. Kyushu Univer.* **56**, 317-326.

- Cohen, A. C., 1965, Maximum likelihood estimation in the Weibull distribution based on complete and on censored samples. *Technometrics*. **7**, 579-588.
- Cordeiro, G. M., Ortega, E.M.M., Silva, G.O., 2014, The Kumaraswamy modified Weibull distribution: theory and applications. *J. Stat. Comput. Simul.* **84**, 1387-1411.
- Dasgupta, R. 2015, Longitudinal growth of elephant foot yam and some characterisation theorems. *Springer Proc. Math. Stat.* **132**, 259-285.
- Dey, S., Dey, T., Anis, M. Z., 2015, Weighted Weibull distribution: Properties and estimation. *J. Stat. Theo. Pract.* **9**, 250-265.
- Elbatal, I., Mansour, M. M., Ahsanullah, M., 2016, The Additive Weibull-Geometric Distribution: Theory and Applications. *J. Stat. Theo. Appl.* **15**, 125-141.
- Elderton, W. P., 1906, Frequency curves and correlation. Charles and Edwin Layton. London.
- Eliopoulos, P.A., Stathas, G.J., Bouras, S.L., 2005, Effects and interactions of temperature, host deprivation and adult feeding on the longevity of the parasitoid *Venturia canescens* (Hymenoptera: Ichneumonidae). *Euro. J. Entomology*. **102**, 181-187.
- Enszer, J. A., Lin, Y., Ferson, S., Corliss, G. F., Stadtherr, M. A., 2011, Probability bounds for nonlinear dynamic process, *AIChE Journal*. **57**, 404-422.
- Fargo, W.S., Wagner, T.L., Coulson, R.N., Cover, J.D., McAudle, T., Schowalter, T.D., 1982, Probability functions for components of the *Dendroctonus frontalis*-Host tree population system and their potential use with population models. *Res. Popul. Ecology*. **24**, 123-131.
- Forrester, D.I., Elms, S.R., Baker, T.G., 2013, Tree growth-competition relationships in thinned Eucalyptus plantations vary with stand structure and site quality. *Euro. J. Forest Res.* **132**, 241-252.
- Gao, D., Perry, G., 2016, Species–area relationships and additive partitioning of diversity of native and nonnative herpetofauna of the West Indies. *Ecology and Evolution*. **6**, 7742-7762.
- Ghitany, M.E., Al-Jarallah, R.A., Balakrishnan, N., 2011, On the existence and uniqueness of the MLEs of the parameters of a general class of exponentiated distributions. *Statistics*. **47**, 37-41.
- Guseinoviene, E., Senulis, A., Akinci, T.C., Seker, S., 2014, Statistical and Continuous Wavelet Analysis of wind speed data in Mardin-Turkey. *9th Int.I Conf. on Ecological Vehicles and Renew. Energies, EVER*. Article number 6844073.
- Hagiwara, Y., 1974, Probability of earthquake occurrence as obtained from a Weibull distribution analysis of crustal strain. *Tectonophysics*. **23**, 313-318.
- Hall, A. J., Gandar, P. W., 1996, Stochastic models for fruit growth. *Acta Horticulture*. **416**, 113-119.

- Hallinan, A. J., 1993, A review of the Weibull distribution. *J. Quality Tech.* **25**, 85-85.
- He, D., Hong, W., Hu, Haiqing, Wu, C., 2004, Patch size distribution pattern and its hierarchical effect of main landscape types in the Wuyishan scenery district. *Chinese J. Appl. Ecology.* **15**, 21-25.
- Hirose, H., 2011, Parameter estimation for the truncated Weibull model using the ordinary differential equation. *Proceedings-1st ACIS/JNU International Conference on Computers, Networks, Systems and Industrial Engineering. CNSI*, 396-399.
- Hirose, H., 2012, Estimation of number of failures in the Weibull model using the ordinary differential equation. *Euro. J. Operat. Res.* **223**, 722-731.
- Hogg, D.B., Nordheim, E.V., 1983, Age-specific survivorship analysis of *Heliothis* spp. Populations on cotton. *Res. Popul. Ecology.* **25**, 280-297.
- Huffman, D.W., Crouse, J.E., Walker Chancellor, W., Fulé, P.Z., 2012, Influence of time since fire on pinyon-juniper woodland structure. *Forest Ecol. Magt.* **274**, 29-37.
- Jankowski, M.D., Williams, C.J., Fair, J.M., Owen, J.C., 2013, Birds Shed RNA-Viruses According to the Pareto Principle. *PLoS ONE.* **8**, Article number e72611.
- Johnson N. L., Kotz, S., Balakrishnan, N., 1994, Continuous Univariate Distributions. Wiley New York. ISBN: 0-471-58495-9.
- Johnson N. L., Kotz, S., Balakrishnan, N., 1995, Continuous Univariate Distributions, Volume 2. 2nd edition, Wiley.
- Kapur, A., Keoleian, G., Kendall, A., Kesler, S.E., 2008, Dynamic modeling of in-use cement stocks in the United States. *J. Indust. Ecology.* **12**, 539-556.
- Knopoff, D., 2013, On the modeling of migration phenomena on small networks. *Math. Mod. Meth. Appl. Sci.* **23**, 541-563.
- Lee, C., Famoye, F., Olumolade, O., 2007, Beta-Weibull distribution: some properties and applications to censored data. *J. Modern Appl. Stat. Meth.* **6**, Article: 17.
- Legge, A.H., Krupa, S.V., 1989, Air quality at a high elevation, remote site in Western Canada. *Proceedings - 82nd A&WMA Annual Meeting; Anaheim, CA, USA; Code 13687, Proceedings - A&WMA Annual Meeting*, **8**, 17.
- Lemon, G. H., 1975, Maximum likelihood estimation for the three parameter Weibull distribution based on censored samples. *Technometrics.* **17**, 247-254.
- Li, W.-C. Wang, S.-D., Zhong, Z.-K., Sheng, H.-Y., Zhou, Y., 2011, Growth rhythm and individual models of wine bamboo plantation, *Oxytenanthera braunii*. *Forest Research.* **24**, 713-719.

- Lieblein, J. 1955, On moments of order statistics from the Weibull distribution. *Ann. Math. Statist.* **26**, 330-333.
- Littell, R. C., Mc Clave, J. T., Offen, W. W., 1979, Goodness-of-fit tests for the two parameter Weibull distribution. *Comm. Stat. Simul. Comput.* **8**, 257-269.
- Luo, Y.-J., Yang, Z.-G., Wang, D., Xie, D.-Y., Ding, W., 2015, Relative fitness of propargite-resistant strain of *Tetranychus cinnabarinus* (Boisduval)(Acari: Tetranychidae). *Chinese J. Ecology.* **34**, 2827-2832.
- MacKenzie, R.A., Kaster, J.L., Klump, J.V., 2004, The ecological patterns of benthic invertebrates in a Great Lakes coastal wetland. *J. Great Lakes Res.* **30**, 58-69.
- Mafart, P., Couvert, O., Gaillard, S., Leguérinel, I., 2002, On calculating sterility in thermal preservation methods: application of the Weibull frequency distribution model. *Int. J. Food Microbio.* **72**, 107-113.
- Marabi, A., Livings, S., Jacobson, M., Saguy, I. S., 2003, Normalized Weibull distribution for modeling rehydration of food particulates. *Euro. Food Res. Technol.* **217**, 311-318.
- McKenzie, D., Hessel, A.E., Kellogg, L.-K.B., 2006, Using neutral models to identify constraints on low-severity fire regimes. *Landscape Ecology*, **21**, 139-152.
- Menon, M. V., 1963, Estimation of the shape and scale parameters of the Weibull distribution. *Technometrics.* **5**, 175-182.
- Mohammadi, K., Alavi, O., Mostafaeipour, A., Goudarzi, N., Jalilvand, M., 2016, Assessing different parameters estimation methods of Weibull distribution to compute wind power density. *Energy Convers. Magt.* **108**, 322-335.
- Moritz, M.A., Moody, T.J., Miles, L.J., Smith, M.M., Valpine, P., 2009, The fire frequency analysis branch of the pyrostatistics tree: Sampling decisions and censoring in fire interval data, *Environ. Ecol. Stat.* **16**, 271-289.
- Mudholkar, G. S., Srivastava, D. K., Kollia, G. D., 1996, A generalization of the Weibull distribution with application to the analysis of survival data. *J. Amer. Stat. Assoc.* **91**, 1575-1583.
- Nadarajah, S., Cordeiro, G. M., Ortega, E. M., 2013, The exponentiated Weibull distribution: a survey. *Stat. Papers.* **54**, 839-877.
- Nakagawa, T., Osaki, S., 1975, The discrete Weibull distribution. *IEEE Transac. Reliability.* **24**, 300-301.

Nanos, N., Montero, G., 2002, Spatial prediction of diameter distribution models. *Forest Ecol. Magt.* **161**, 147-158.

Nassar, M. M., Eissa, F. H., 2003, On the exponentiated Weibull distribution. *Comm. Stat. Theo. Meth.* **32**, 1317-1336.

Navarro-Cerrillo, R.M., Olave, F., Moreno, F., de Miguel, S., Clemente, M., 2014, Stand structure and regeneration of harvested *Araucaria araucana*-*Nothofagus* stands in central Chile, *Southern Forests.* **76**, 11-19.

Neeff, T., Dutra, L.V., Dos Santos, J.R., Da Costa Freitas, C., Araujo, L.S. 2003, Tropical forest stand table modelling from SAR data. *Forest Ecol. Magt.* **186**, 159-170.

Newman, M.C., Aplin, M.S., 1992, Enhancing toxicity data interpretation and prediction of ecological risk with survival time modeling: an illustration using sodium chloride toxicity to mosquitofish (*Gambusia holbrooki*). *Aquatic Toxicology.* **23**, 85-96.

Oguntunde, P. E., Balogun, O. S., Okagbue, H. I., Bishop, S. A., 2015, The Weibull-Exponential Distribution: Its Properties and Applications. *J. Appl. Sci.* **15**, 1305-1311.

Okagbue, H.I., Oguntunde, P.E., Opanuga, A.A., Owoloko, E.A. "Classes of Ordinary Differential Equations Obtained for the Probability Functions of Fréchet Distribution," *Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 186-191.*

Okagbue, H.I., Oguntunde, P.E., Ugwoke, P.O., Opanuga, A.A. "Classes of Ordinary Differential Equations Obtained for the Probability Functions of Exponentiated Generalized Exponential Distribution," *Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 192-197.*

Okagbue, H.I., Opanuga, A.A., Owoloko, E.A., Adamu, M.O. "Classes of Ordinary Differential Equations Obtained for the Probability Functions of Cauchy, Standard Cauchy and Log-Cauchy Distributions," *Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 198-204.*

Okagbue, H.I., Bishop, S.A., Opanuga, A.A., Adamu, M.O. "Classes of Ordinary Differential Equations Obtained for the Probability Functions of Burr XII and Pareto Distributions," *Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 399-404.*

Okagbue, H.I., Adamu, M.O., Owoloko, E.A., Opanuga, E.A. "Classes of Ordinary Differential Equations Obtained for the Probability Functions of Gompertz and Gamma Gompertz Distributions," Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 405-411.

Okagbue, H.I., Adamu, M.O., Anake, T.A. "Quantile Approximation of the Chi-square Distribution using the Quantile Mechanics," Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 477-483.

Okagbue, H.I., Adamu, M.O., Opanuga, A.A., Oghonyon, J.G. "Classes of Ordinary Differential Equations Obtained for the Probability Functions of 3-Parameter Weibull Distribution," Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 539-545.

Okagbue, H.I., Opanuga, A.A., Owoloko, E.A., Adamu, M.O. "Classes of Ordinary Differential Equations Obtained for the Probability Functions of Exponentiated Fréchet Distribution," Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 546-551.

Okagbue, H.I., Adamu, M.O., Owoloko, E.A., Bishop, S.A. "Classes of Ordinary Differential Equations Obtained for the Probability Functions of Half-Cauchy and Power Cauchy Distributions," Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 552-558.

Okagbue, H.I., Adamu, M.O., Anake, T.A. "Solutions of Chi-square Quantile Differential Equation," Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 813-818.

Okagbue, H.I., Oguntunde, P.E., Opanuga, A.A., Owoloko, E.A. "Classes of Ordinary Differential Equations Obtained for the Probability Functions of Exponential and Truncated Exponential Distributions," Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 858-864.

Okagbue, H.I., Agboola, O.O., Ugwoke, P.O., Opanuga, A.A. "Classes of Ordinary Differential Equations Obtained for the Probability Functions of Exponentiated Pareto Distribution," Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 865-870.

Okagbue, H.I., Agboola, O.O., Opanuga, A.A., Oghonyon, J.G. "Classes of Ordinary Differential Equations Obtained for the Probability Functions of Gumbel Distribution," Lecture Notes in

Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 871-875.

Okagbue, H.I., Odetunmibi, O.A., Opanuga, A.A., Oguntunde, P.E. "Classes of Ordinary Differential Equations Obtained for the Probability Functions of Half-Normal Distribution," Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 876-882.

Okagbue, H.I., Adamu, M.O., Owoloko, E.A., Suleiman, E.A. "Classes of Ordinary Differential Equations Obtained for the Probability Functions of Harris Extended Exponential Distribution," Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering and Computer Science 2017, 25-27 October, 2017, San Francisco, U.S.A., pp 883-888.

Papadimitriou, Katafygiotis, L. S., Beck, J. L., 1995, Approximate analysis of response variability of uncertain linear systems. *Probab. Engineering Mech.* **10**, 251-264.

Percontini, A., Blas, B., Cordeiro, G. M., 2013, The beta Weibull Poisson distribution. *Chilean J. Stat.* **4**, 3-26.

Pham, H., Lai, C. D., 2007, On recent generalizations of the Weibull distribution. *IEEE Transac. Reliability*, **56**, 454-458.

Pinder, J. E., Wiener, J. G., Smith, M. H., 1978, The Weibull distribution: a new method of summarizing survivorship data. *Ecology*, **59**, 175-179.

Pogány, T. K., Saboor, A., Provost, S., 2015, The Marshall–Olkin exponential Weibull distribution. *Hacettepe J. Math. Stat.* **14**, 1579-1594.

Razali, A. M., Al-Wakeel, A. A., 2013, Mixture Weibull distributions for fitting failure times data. *Appl. Math. Comput.* **219**, 11358-11364.

Reed, W. J., Hughes, B. D., 2002, On the size distribution of live genera. *J. Theo. Biology.* **217**, 125-135.

Rinne H., 2010, Location scale Distributions, Linear Estimation and probability plotting using MATLA

Rogeanu, M.-P., Armstrong, G.W., 2017, Quantifying the effect of elevation and aspect on fire return intervals in the Canadian Rocky Mountains. *Forest Ecol. Magt.* **384**, 248-261.

Roylance, B. J., Pocock, G., 1983, Wear studies through particle size distribution I: Application of the Weibull distribution to ferrography. *Wear*, **90**, 113-136.

Saboor, A., Pogany, T. K., 2016, Marshall-Olkin gamma-Weibull distribution with applications. *Comm. Stat. Theo. Meth.* **45**, 1550-1563.

- Saboor, A., Kamal, M., Ahmad, M., 2015, The transmuted exponential–Weibull distribution with applications. *Pak. J. Stat.* **31**, 229-250.
- Sarkkola, S., Alenius, V., Hökkä, H., Laiho, R., Päivänen, J., Penttilä, T., 2003, Changes in structural inequality in Norway spruce stands on peatland sites after water-level drawdown. *Canadian J. Forest Res.* **33**, 222-231.
- Sarkkola, S., Hökkä, H., Laiho, R., Päivänen, J., Penttilä, T., 2005, Stand structural dynamics on drained peatlands dominated by Scots pine. *Forest Ecol. Magt.* **206**, 135-152.
- Sedaghat, A. , Gaith, M., Khanafer, K., Bani-Hani, E., 2016, Rated wind speed reality or myth for optimization in design of wind turbines. *11th Int. Conf. on Ecol. Vehicles and Renew. Energies, EVER*. Article number 7476402.
- Seguro, J. V., Lambert, T. W., 2000, Modern estimation of the parameters of the Weibull wind speed distribution for wind energy analysis. *J. Wind Enginee. Indust. Aerody.* **85**, 75-84.
- Selim, M. A., Badr, A. M., 2016, Kumaraswamy Generalized Power Weibull Distribution. *Math. Theo. Model.*, **6**, 110-124.
- Sharif, M. N., Islam, M. N., 1980, The Weibull distribution as a general model for forecasting technological change. *Technol. Forecast. Social Change.* **18**, 247-256.
- Shittu, O. I., Adepoju, K. A., 2014, On the Exponentiated Weibull Distribution for Modeling Wind Speed in South Western Nigeria. *J. Modern Appl. Stat. Meth.* **13**, Article: 28.
- Shotorban, B., 2010, Dynamic least-square kernel density modeling of Fokker-Planck equations with application to neural population. *Physical Review E-Stat. Nonl. Soft Matt. Phys.* **81**, Article number: 046706.
- Singh, V. P., 1987, On application of the Weibull distribution in hydrology. *Water Resource Magt.* **1**, 33-43.
- Singla, N., Jain, K., Sharma, S. K., 2012, The beta generalized Weibull distribution: properties and applications. *Relia. Engine. Syst. Safety*, **102**, 5-15.
- Stauffer, H. B., 1979, A derivation for the Weibull distribution. *J. Theo. Biology.* **81**, 55-63.
- Steinsaltz, D., Evans, S. N., Wachter, K. W., 2005, A generalized model of mutation-selection balance applications to aging. *Adv. Appl. Math.* **35**, 16-33.
- Studds, C.E., McFarland, K.P., Aubry, Y., Rimmer, C.C., Hobson, K.A., Marra, P.P., Wassenaar, L.I., 2012, Stable-hydrogen isotope measures of natal dispersal reflect observed population declines in a threatened migratory songbird. *Divers. Distribut.* **18**, 919-930.

- Subedi, S., Fox, T.R., 2016, Modeling repeated fertilizer response and one-time midrotation fertilizer response in loblolly pine plantations using FR in the 3-PG process model. *Forest Ecol. Magt.* **380**, 90-99.
- Tadikamalla, P. R., 1978, Applications of the Weibull distribution in inventory control. *J. Operational Res. Soc.* **29**, 77-83.
- Tahir, M. H., Cordeiro, G. M., Alzaatreh, A., Mansoor, M., Zubair, M. (2016 b). A New Weibull–Pareto Distribution: Properties and Applications. *Comm. Stat. Simul. Comput.* **45**, 3548-3567.
- Tahir, M. H., Cordeiro, G. M., Mansoor, M., Zubair, M., Alizadeh, M., 2016 c, The Weibull–Dagum distribution: Properties and applications. *Comm. Stat. Theo. Meth.* **45**, 7376-7398.
- Tahir, M. H., Zubair, M., Mansoor, M., Cordeiro, G. M., Alizadeh, M., Hamedani, G. G., 2016 d, A new Weibull-G family of distributions. *Hacettepe J. Math. Stat.* **45**, 629-647.
- Tahir, M., Alizadeh, M., Mansoor, M., Cordeiro, G. M., Zubair, M., 2016 a, The Weibull-power function distribution with applications. *Hacettepe J. Math. Stat.* **45**, 245-265.
- Tahir, M.H., Cordeiro, G.M., Mansoor, M., Zubair, M., 2015, The Weibull-Lomax distribution: properties and applications. *Hacettepe J. Math. Stat.* **44**, 455-474.
- Taubert, F., Hartig, F., Dobner, H.-J., Huth, A., 2013, On the Challenge of Fitting Tree Size Distributions in Ecology. *PLoS ONE*, **8**, Article number e58036.
- Thoman, D. R., Bain, L. J., Antle, C. E., 1969, Inferences on the parameters of the Weibull distribution. *Technometrics.* **11**, 445-460.
- Tjørve, E., 2003, Shapes and functions of species-area curves: A review of possible models. *J. Biogeography.* **30**, 827-835.
- Wagner, T. L., Wu, H. I., Sharpe, P. J., Coulson, R. N., 1984, Modeling distributions of insect development time: a literature review and application of the Weibull function. *Ann. Entomol. Soc. Amer.* **77**, 475-483.
- Wang, D.-H. , Hsieh, H.-C., Tang, S.-C., Chung, C.-H., 2010, Stand growth simulation of a Taiwania plantation in the Liouguei area. *Taiwan J. Forest Sci.* **25**, 155-169.
- Wong, T. F., Wong, R. H., Chau, K. T., Tang, C. A., 2006, Microcrack statistics, Weibull distribution and micromechanical modeling of compressive failure in rock. *Mechanics of Materials.* **38**, 664-681.
- Yang, J., He, H.S., Gustafson, E.J., 2004, A hierarchical fire frequency model to simulate temporal patterns of fire regimes in LANDIS. *Ecolog. Modell.* **180**, 119-133.

Zabel, R.W., Burke, B.J., Moser, M.L., Caudill, C.C., 2014, Modeling temporal phenomena in variable environments with parametric models: AN application to migrating salmon. *Ecolog. Model.* **273**, 23-30.